

Understanding HIV infection

Fifth Edition

Terrence  
HIGGINS  
TRUST



# HIV? AIDS?



This booklet is for you if you have any basic questions about HIV: what it is, how it is transmitted, and its effects on the human body. If you have HIV infection and have concerns about anti-HIV treatment, side-effects and how to live well with HIV, Terrence Higgins Trust produce a range of information that could be useful.

This leaflet covers:

- 1: What is HIV and AIDS?
- 2: How is HIV transmitted?
- 3: What is the HIV test?
- 4: How does HIV affect the body?
- 5: Living with HIV

# 1: What are HIV and AIDS?

**HIV** stands for **Human Immunodeficiency Virus**.

It is a viral infection that attacks the human body's immune system. Initially a person may show no symptoms of HIV infection as their immune system manages to control it. However, in most cases, the immune system will require help from anti-HIV drugs to keep the HIV infection under control. These drugs do not completely rid the body of HIV infection.

**AIDS (Acquired Immune Deficiency Syndrome)**

as a term is now very rarely used. It is more usual to talk of late-stage or advanced HIV infection. AIDS is not a single disease or condition, it is a term that is used to describe the point when a person's immune system can no longer cope, because of damage caused by HIV and specific illnesses start to appear. People do not actually die from AIDS; they die from the cancers, pneumonia or other conditions that may take hold when their immune system has been weakened by HIV.

## 2: How is HIV transmitted and how can the risks be reduced?

As far as viruses go, HIV is not an easy one to pass from one person to another. The virus has to get out of an HIV infected person's body and into the blood stream of someone who does not have HIV, for that person to become infected.

Body fluids which contain enough HIV to infect someone are:

- sperm and seminal fluid
- vaginal fluids, including menstrual fluids
- breast milk
- blood

Other body fluids like saliva, sweat or urine, do not contain enough of the virus to infect another person.

The following is a list of possible ways that HIV can be passed from one person to another, and information on how the risk can be reduced. It's important to remember that at least one of the people has to have HIV for there to be any risk of infection.

## Semen, seminal fluid, 'cum'

If an HIV infected man's penis enters an uninfected person's body there could be HIV infection present in the 'pre cum' (which is the fluid released from the penis to help lubricate it). HIV is also found in much greater quantities in the semen and can be passed on through vaginal, anal and oral sex.

### Reducing the risk

When having penetrative sex (this means that a man's penis is put into someone else), the most effective way of reducing the risk of HIV being passed from one person to another is to use a condom properly with the right lubricant, if needed. It's also important to put the condom on fully before penetration and to check that it stays on the penis. Relying on the man to withdraw before ejaculation has been shown **not** to be an effective method of preventing HIV infection.

Anal and vaginal sex are the most common ways that HIV is passed from someone with HIV to another person. There have been some cases of HIV being passed on during oral sex when someone who doesn't have HIV sucks the penis of an HIV positive man. This relatively small risk can be reduced by using condoms for oral sex. Some people find flavoured condoms more pleasant for oral sex.

HIV is not usually passed on when:

- a man who does not have HIV receives oral sex (has it done to him) from someone who is HIV positive
- a woman receives oral sex (has it done to her) from someone who is HIV positive

### **Mother to baby** (sometimes called vertical transmission)

If a woman with HIV becomes pregnant, there is a risk of her passing on HIV infection to her baby during birth. While in the womb, the baby remains fairly protected against HIV infection, but during the trauma of birth HIV can be passed on. HIV may also be transmitted if the HIV positive woman breastfeeds her baby, as HIV has been detected in the breast-milk of women with HIV and the digestive systems of young babies are not fully developed, making them more at risk of infection.

### **Reducing the risk**

For a woman with HIV it is possible for her child to be born without HIV. This can be achieved through careful planning by the woman and medical professionals, and by using certain anti-HIV drugs at particular points during the pregnancy. Mothers with HIV are also advised to bottle-feed their babies rather than breastfeed to reduce the risk of HIV being passed through breast-milk.

## Blood to blood

Bleeding on its own isn't enough to pass on HIV infection. The blood of someone with HIV needs to leave their body and get into an uninfected person. Before 1985/6 a number of people, including some with haemophilia, were infected through blood products and blood transfusions. Since 1985/6 all blood donations have been screened for HIV and blood products have been treated.

Another way that blood can be passed from one person to another is through sharing needles and this has led to some intravenous drug users being infected with HIV.

### Reducing the risk

For intravenous drug users (those who inject into the body, which includes people who inject steroids into their muscles), the main ways that the risk of HIV infection can be reduced are to always use fresh needles and never share drug equipment with anyone else.

HIV is not passed on from one person to another if:

- Neither person is HIV positive. Neither a person's body or having sex creates HIV: one of the people involved has to be infected with HIV already.
- An uninfected person touches or is close to someone with HIV.
- HIV positive and uninfected people share cups, glasses and cutlery.

### Post Exposure Prophylaxis (PEP)

PEP is a course of anti-HIV drugs which may prevent HIV infection after HIV has entered the body.

Treatment must be started as soon as possible after the potential exposure to HIV.

PEP treatment is available through local sexual health clinics or hospital Accident and Emergency (A&E) departments. For up-to-date information go to [www.tht.org.uk/pep](http://www.tht.org.uk/pep).

## 3: What is the HIV test?

The HIV test shows whether someone is infected with HIV. It is not a test for AIDS.

### Antibodies

The most commonly available HIV test is the HIV antibody test, which looks for antibodies to HIV. These antibodies are created by the infected person's immune system in response to the presence of HIV.

### The window period (seroconversion)

When someone becomes infected with HIV, it can take up to three months for their immune system to produce enough antibodies to show up on an HIV test. This gap is called the *window period*, or *seroconversion*. Throughout this period, the person with HIV has enough of the virus in their blood, sexual fluids and breast-milk to infect another person. Often the level of virus in their body fluids at this time can be extremely high, making them particularly infectious.

Because the HIV test looks for antibodies, taking an HIV test less than three months after possible infection might not give an accurate result.

### A positive result

If the test finds HIV antibodies, the result is said to be positive. The person would then be HIV antibody positive or 'HIV positive'. In other words this person has HIV. It does not mean that they have AIDS. The test does not show if, when or whether they will become ill. But it can be used as a guide as to whether the time is right to start anti-HIV treatment.

### A negative result

If the test does not find HIV antibodies, the result is said to be negative. A negative test result shows that the person does not have detectable HIV infection at this time. This is accurate as long as the test was done at the end of the 'window period'.

## 4: How does HIV affect the body?

After a while, HIV will begin to weaken the body's immune system, which usually fights off any infections.

HIV reproduces even in someone who may be feeling well. If their immune system is working well, the virus may only reproduce a little. But if their immune system is weakened through illness, the virus can reproduce itself much more easily. Research shows that people with higher levels of HIV in their bodies have a greater risk of becoming ill than people with less HIV in their bodies.

As the HIV reproduces itself, the person is more vulnerable to other infections. Normally many of these infections would be controlled by an effective immune system, but in someone with HIV and a weakened immune system they can become more serious and in some cases fatal. These illnesses are often called 'opportunistic infections'.

Using anti-HIV treatments can slow this process down. They do not clear the body of HIV infection but do help control the amount of virus. If the HIV is controlled, the person's immune system is better able to deal with any other infections.

### How effective is anti-HIV treatment?

Taking a combination of anti-HIV drugs (combination therapy) can successfully reduce the level of HIV in the blood, and in many cases have a dramatic effect in improving the health and life expectancy of someone with HIV. However, the longer term effects of these treatments and for how long they remain effective is still unknown.

### Is there now a cure for HIV?

At present there is no cure for HIV infection. Drug combinations do not completely eliminate HIV from the body; they at best control it. There is still no vaccine to prevent people becoming infected with HIV.

## 5: Living with HIV

Many people manage to continue living with HIV without many alterations to their lives; remaining in work, relationships, raising or starting families and forming new relationships. The main aspect of life that can sometimes dramatically alter is other people's attitudes towards them if they know or find out that they have HIV. Other people's attitudes can range from being supportive and accepting, through to

rejection and hostility. Because of this, the hardest part of life with HIV for many people is telling others, as they are uncertain of what their reaction will be. Some people have been rejected by family, friends and partners.

As a result of this many HIV positive people choose not to tell anyone about being HIV positive except for a few trusted friends. Not being able to talk openly about concerns can be hard, but there are often support groups around the country available to help counter this feeling of isolation.

**For further information and details of these support groups call THT DIRECT > 0845 12 21 200**

## Prosecutions

There have been a few cases of people with HIV being prosecuted for transmitting HIV to their sexual partners. Most HIV positive people make a considerable effort to avoid infecting others. However, many people with HIV do not know they have it, so cannot tell the people that have sex with. Condoms provide best protection against infection. For up-to-date information, go to [www.tht.org.uk/prosecutions](http://www.tht.org.uk/prosecutions)

## Terrence Higgins Trust Offices

We are opening offices all the time. These details were correct when this leaflet was printed in June 2007. Please check our website [www.tht.org.uk](http://www.tht.org.uk) for up-to-date information on our locations throughout the United Kingdom.

### England Offices

Terrence Higgins Trust  
Email: [info@tht.org.uk](mailto:info@tht.org.uk)

#### Birkenhead

New office opening soon  
Contact THT Direct 0845 1221 200  
for more details or email  
[info.wirral@tht.org.uk](mailto:info.wirral@tht.org.uk)

#### Birmingham

Terrence Higgins Trust,  
Unit G7, The Arch,  
48-52 Floodgate Street,  
Birmingham, B5 5SL  
Tel: 0121 694 6440 Fax: 0121 694 6441  
Email: [info.birmingham@tht.org.uk](mailto:info.birmingham@tht.org.uk)

#### Brighton

Terrence Higgins Trust,  
61 Ship Street, Brighton,  
East Sussex, BN1 1AE  
Tel: 01273 764 200 Fax: 01273 764 201  
Email: [info.brighton@tht.org.uk](mailto:info.brighton@tht.org.uk)

#### Bristol

Terrence Higgins Trust,  
Aled Richards Centre,  
8-10 West Street, Old Market,  
Bristol, BS2 0BH  
Tel: 0117 955 1000 Fax: 0117 954 1200  
Email: [info.bristol@tht.org.uk](mailto:info.bristol@tht.org.uk)

#### Colchester

Terrence Higgins Trust,  
The Old Customs House,  
Hythe Quay, Colchester,  
Essex, CO2 8JB  
Tel: 01206 798 595 Fax: 01206 792 934  
Email: [info.colchester@tht.org.uk](mailto:info.colchester@tht.org.uk)

#### Coventry

Terrence Higgins Trust,  
10 Manor Road,  
Coventry, CV1 2LH  
Tel: 02476 229 292 Fax: 02476 634 248  
Email: [info.coventry@tht.org.uk](mailto:info.coventry@tht.org.uk)

#### Eastbourne

Terrence Higgins Trust,  
Rooms 9-14 Dyke House,  
110 South Street, Eastbourne,  
East Sussex, BN21 4LZ  
Tel: 01323 649 927 Fax: 01323 439 340  
Email: [info.eastbourne@tht.org.uk](mailto:info.eastbourne@tht.org.uk)

#### London - Camberwell

Terrence Higgins Trust,  
Lighthouse King's, Unit 2, Empress Mews  
Kenbury Street, Camberwell  
London, SE5 9BT  
Tel: 020 7737 9740 Fax: 020 7737 9741  
Email [info.camberwell@tht.org.uk](mailto:info.camberwell@tht.org.uk)

#### London - Counselling

Terrence Higgins Trust,  
314-320 Gray's Inn Road,  
London, WC1X 8DP  
Tel: 020 7812 1777 Fax: 020 7812 1742  
Email: [info.counselling@tht.org.uk](mailto:info.counselling@tht.org.uk)

#### London - Earls Court

Terrence Higgins Trust,  
SW5, 11 Eardley Crescent,  
London, SW5 9JS  
Tel: 020 7370 0406 Fax: 020 7244 0037  
Email: [info.sw5@tht.org.uk](mailto:info.sw5@tht.org.uk)

#### London - Ladbroke Grove

Terrence Higgins Trust,  
Lighthouse West London,  
111-117 Lancaster Road,  
London, W11 1QT  
Tel: 020 7313 2900 Fax: 020 7229 1258  
Email: [info.ladbrokegrove@tht.org.uk](mailto:info.ladbrokegrove@tht.org.uk)

#### Leeds

New office opening soon  
Contact THT Direct 0845 1221 200  
for more details or email  
[info.leeds@tht.org.uk](mailto:info.leeds@tht.org.uk)

#### London - Kings Cross

Terrence Higgins Trust,  
314-320 Gray's Inn Road,  
London, WC1X 8DP  
Tel: 020 7812 1600 Fax: 020 7812 1601  
Email: [info.@tht.org.uk](mailto:info.@tht.org.uk)

### **London – London Bridge**

Terrence Higgins Trust,  
Healthy Gay Living Counselling,  
Hop Exchange, Unit 34, 1st Floor,  
24 Southwark Street, London, SE1 1TY  
Tel: 020 7378 6386  
Email: [info.counselling@tht.org.uk](mailto:info.counselling@tht.org.uk)

### **London – Waterloo**

Terrence Higgins Trust,  
Lighthouse South London,  
14-15 Lower Marsh,  
London, SE1 7RJ  
Tel: 020 7803 1660 Fax: 020 7803 1661

### **Oxford**

Terrence Higgins Trust,  
43 Pembroke Street,  
Oxford, OX1 1BP  
Tel: 01865 243 389 Fax: 01865 792 210  
Email: [info.oxford@tht.org.uk](mailto:info.oxford@tht.org.uk)

### **Southend-on-Sea**

Terrence Higgins Trust,  
20 Weston Road, Southend-on-Sea,  
Essex, SS1 1AS  
Tel: 01702 340 791 Fax: 01702 335 766  
Email: [info.southend@tht.org.uk](mailto:info.southend@tht.org.uk)

### **Swindon**

Terrence Higgins Trust,  
Shaftesbury Centre,  
Percy Street, Swindon,  
Wiltshire, SN2 2AZ.  
Tel: 01793 554 825  
Email: [info.swindon@tht.org.uk](mailto:info.swindon@tht.org.uk)

### **Telford**

Terrence Higgins Trust,  
4 Park Street, Wellington, Telford,  
Shropshire, TF1 3AE  
Tel: 01952 221 410 Fax: 01952 221 420  
Email: [info.shropshire@tht.org.uk](mailto:info.shropshire@tht.org.uk)

### **Woking**

Terrence Higgins Trust,  
Regent House, 19-20 The Broadway,  
Woking, Surrey, GU21 5XQ  
Tel: 01483 263 160 Fax: 01483 263 170  
Email: [info.woking@tht.org.uk](mailto:info.woking@tht.org.uk)

### **Wolverhampton**

Terrence Higgins Trust,  
242-244 Bond House, Bond Street,  
Wolverhampton, WV2 4AS  
Tel: 01902 711 818 Fax: 01902 711 925  
Email: [info.wolverhampton@tht.org.uk](mailto:info.wolverhampton@tht.org.uk)

## **Scotland Offices**

Terrence Higgins Trust Scotland  
Email: [info.scotland@tht.org.uk](mailto:info.scotland@tht.org.uk)

### **Aberdeen**

Terrence Higgins Trust Scotland,  
11 Waverley Place,  
Aberdeen, AB10 1XH  
Tel: 0845 241 2151 Fax: 0845 241 2152  
Email: [info.aberdeen@tht.org.uk](mailto:info.aberdeen@tht.org.uk)

### **Glasgow**

Terrence Higgins Trust Scotland,  
Top Floor, Rothesay House,  
134 Douglas Street, Glasgow G2 4HF  
Tel: 0141 332 3838 Fax: 0141 332 3755  
Email: [info.scotland@tht.org.uk](mailto:info.scotland@tht.org.uk)

### **Inverness**

Terrence Higgins Trust Scotland,  
34 Waterloo Place,  
Inverness IV1 1NB  
Tel: 01463 711 585 Fax: 01463 711 793  
Email: [info.highland@tht.org.uk](mailto:info.highland@tht.org.uk)

## **Wales Offices**

Terrence Higgins Trust Cymru  
Email: [info.cymru@tht.org.uk](mailto:info.cymru@tht.org.uk)

### **Cardiff**

Terrence Higgins Trust Cymru,  
Canton House,  
435-451 Cowbridge Road East,  
Canton, Cardiff, CF5 1JH  
Tel/Fax: 029 2066 6465  
Email: [info.cymru@tht.org.uk](mailto:info.cymru@tht.org.uk)

### **Swansea**

Terrence Higgins Trust Cymru,  
c/o YMCA, 1 Kingsway, Swansea, SA1 5JQ  
Tel/Fax: 01792 477 540  
Email: [info.cymru@tht.org.uk](mailto:info.cymru@tht.org.uk)

# The HIV and sexual health charity for life

Website: [www.tht.org.uk](http://www.tht.org.uk) THT Direct: 0845 12 21 200

Registered office: 314-320 Gray's Inn Road, London WC1X 8DP

Tel: 020 7812 1600 Email: [info@tht.org.uk](mailto:info@tht.org.uk)

© Terrence Higgins Trust, June 2007 Code: 212

Terrence Higgins Trust is a registered charity no. 288527. Company reg. no.1778149.

Registered in England. A company limited by guarantee.

Design Felton Communication 020 7405 0900 Ref: 9865.2